Auto2 prover

Bohua Zhan

April 18, 2024

Abstract

Auto2 is a saturation-based heuristic prover for higher-order logic, implemented as a tactic in Isabelle.

This entry contains the instantiation of auto2 for Isabelle/HOL, along with two basic examples: solutions to some of the Pelletier's problems, and elementary number theory of primes.

Contents

1 Introduction **3**

2 Pelletier's problems **3**

3 Primes **6**

3.1 Basic definition 7

3.2 Infinitude of primes 8

3.3 Existence and uniqueness of prime factorization 8

1 Introduction

Auto2 [2] is a proof automation tool implemented in Isabelle. It uses a saturation-based approach to proof search: starting with a list of initial assumptions, it iteratively adds facts that can be derived from these assumptions, with the aim of ultimately deriving a contradiction. Users can add their own proof procedures to auto2 in the form of *proof steps*, in order to implement domain-specific knowledge. Auto2 can be instantiated to both Isabelle/HOL (for ordinary usage) and Isabelle/FOL (for formalization of mathematics based on set theory).

This AFP entry contains the instantiation of auto2 to Isabelle/HOL, and two basic applications:

- Pelletier's problems: solutions to some of the problems in Pelletier's collection of problems for testing automatic theorem provers [1]. Auto2 is not intended to compete with ATPs. In our examples, we merely show how to use the prover to solve some of the problems, sometimes with hints.
- Elementary number theory: theory of prime numbers up to the infinitude of primes and unique factorization. This example follows the development in HOL/Computational_Algebra/Primes.thy in the Isabelle distribution.

2 Pelletier's problems

theory *Pelletier*
  imports *Logic-Thms*
begin

theorem *p1*: $(p \longrightarrow q) \longleftrightarrow (\neg q \longrightarrow \neg p)$ ⟨*proof*⟩

theorem *p2*: $(\neg\neg p) \longleftrightarrow p$ ⟨*proof*⟩

theorem *p3*: $\neg(p \longrightarrow q) \Longrightarrow q \longrightarrow p$ ⟨*proof*⟩

theorem *p4*: $(\neg p \longrightarrow q) \longleftrightarrow (\neg q \longrightarrow p)$ ⟨*proof*⟩

theorem *p5*: $(p \vee q) \longrightarrow (p \vee r) \Longrightarrow p \vee (q \longrightarrow r)$ ⟨*proof*⟩

theorem *p6*: $p \vee \neg p$ ⟨*proof*⟩

theorem *p7*: $p \vee \neg\neg\neg p$ ⟨*proof*⟩

theorem *p8*: $((p \longrightarrow q) \longrightarrow p) \Longrightarrow p$ ⟨*proof*⟩

theorem *p9*: $(p \vee q) \wedge (\neg p \vee q) \wedge (p \vee \neg q) \Longrightarrow \neg(\neg p \vee \neg q)$ ⟨*proof*⟩

theorem *p10*: $q \longrightarrow r \Longrightarrow r \longrightarrow p \wedge q \Longrightarrow p \longrightarrow q \vee r \Longrightarrow p \longleftrightarrow q$ ⟨*proof*⟩

theorem *p11*: $p \longleftrightarrow p$ ⟨*proof*⟩

theorem *p12*: $((p \longleftrightarrow q) \longleftrightarrow r) \longleftrightarrow (p \longleftrightarrow (q \longleftrightarrow r))$
⟨*proof*⟩

theorem *p13*: $p \vee (q \wedge r) \longleftrightarrow (p \vee q) \wedge (p \vee r)$ ⟨*proof*⟩

theorem *p14*: $(p \longleftrightarrow q) \longleftrightarrow ((q \vee \neg p) \wedge (\neg q \vee p))$ ⟨*proof*⟩

theorem *p15*: $(p \longrightarrow q) \longleftrightarrow (\neg p \vee q)$ ⟨*proof*⟩

theorem *p16*: $(p \longrightarrow q) \vee (q \longrightarrow p)$ ⟨*proof*⟩

theorem *p17*: $(p \wedge (q \longrightarrow r) \longrightarrow s) \longleftrightarrow (\neg p \vee q \vee s) \wedge (\neg p \vee \neg r \vee s)$ ⟨*proof*⟩

theorem *p18*: $\exists\, y{::}'a.\ \forall\, x.\ F(y) \longrightarrow F(x)$
⟨*proof*⟩

theorem *p19*: $\exists\, x{::}'a.\ \forall\, y\ z.\ (P(y) \longrightarrow Q(z)) \longrightarrow (P(x) \longrightarrow Q(x))$
⟨*proof*⟩

theorem *p20*: $\forall\, x\ y.\ \exists\, z.\ \forall\, w.\ P(x) \wedge Q(y) \longrightarrow R(z) \wedge S(w) \Longrightarrow$
$\exists\, x\ y.\ P(x) \wedge Q(y) \Longrightarrow \exists\, z.\ R(z)$
⟨*proof*⟩

theorem *p21*: $\exists\, x.\ p \longrightarrow F(x) \Longrightarrow \exists\, x.\ F(x) \longrightarrow p \Longrightarrow \exists\, x.\ p \longleftrightarrow F(x)$
⟨*proof*⟩

theorem *p22*: $\forall\, x{::}'a.\ p \longleftrightarrow F(x) \Longrightarrow p \longleftrightarrow (\forall\, x.\ F(x))$
⟨*proof*⟩

theorem *p23*: $(\forall\, x{::}'a.\ p \vee F(x)) \longleftrightarrow (p \vee (\forall\, x.\ F(x)))$ ⟨*proof*⟩

theorem *p29*: $\exists\, x.\ F(x) \Longrightarrow \exists\, x.\ G(x) \Longrightarrow$
$((\forall\, x.\ F(x) \longrightarrow H(x)) \wedge (\forall\, x.\ G(x) \longrightarrow J(x))) \longleftrightarrow$
$(\forall\, x\ y.\ F(x) \wedge G(y) \longrightarrow H(x) \wedge J(y))$
⟨*proof*⟩

theorem *p30*: $\forall\, x.\ F(x) \vee G(x) \longrightarrow \neg H(x) \Longrightarrow$
$\forall\, x.\ (G(x) \longrightarrow \neg I(x)) \longrightarrow F(x) \wedge H(x) \Longrightarrow \forall\, x.\ I(x)$
⟨*proof*⟩

theorem *p31*: $\neg(\exists\, x.\ F(x) \wedge (G(x) \vee H(x))) \Longrightarrow \exists\, x.\ I(x) \wedge F(x) \Longrightarrow \forall\, x.\ \neg H(x)$
$\longrightarrow J(x) \Longrightarrow$
$\exists\, x.\ I(x) \wedge J(x)$ ⟨*proof*⟩

theorem *p32*: $\forall x.\ (F(x) \wedge (G(x) \vee H(x))) \longrightarrow I(x) \Longrightarrow \forall x.\ I(x) \wedge H(x) \longrightarrow J(x) \Longrightarrow$
$\forall x.\ K(x) \longrightarrow H(x) \Longrightarrow \forall x.\ F(x) \wedge K(x) \longrightarrow J(x)$ ⟨*proof*⟩

theorem *p33*: $(\forall x.\ p(a) \wedge (p(x) \longrightarrow p(b)) \longrightarrow p(c)) \longleftrightarrow$
$(\forall x.\ (\neg p(a) \vee p(x) \vee p(c)) \wedge (\neg p(a) \vee \neg p(b) \vee p(c)))$ ⟨*proof*⟩

theorem *p35*: $\exists (x::'a)\ (y::'b).\ P(x,y) \longrightarrow (\forall x\ y.\ P(x,y))$ ⟨*proof*⟩

theorem *p39*: $\neg(\exists x.\ \forall y.\ F(y,x) \longleftrightarrow \neg F(y,y))$
⟨*proof*⟩

theorem *p40*: $\exists y.\ \forall x.\ F(x,y) \longleftrightarrow F(x,x) \Longrightarrow \neg(\forall x.\ \exists y.\ \forall z.\ F(z,y) \longleftrightarrow \neg F(z,x))$
⟨*proof*⟩

theorem *p42*: $\neg(\exists y.\ \forall x.\ F(x,y) \longleftrightarrow \neg(\exists z.\ F(x,z) \wedge F(z,x)))$
⟨*proof*⟩

theorem *p43*: $\forall x\ y.\ Q(x,y) \longleftrightarrow (\forall z.\ F(z,x) \longleftrightarrow F(z,y)) \Longrightarrow$
$\forall x\ y.\ Q(x,y) \longleftrightarrow Q(y,x)$ ⟨*proof*⟩

theorem *p47*:
$(\forall x.\ P1(x) \longrightarrow P0(x)) \wedge (\exists x.\ P1(x)) \Longrightarrow$
$(\forall x.\ P2(x) \longrightarrow P0(x)) \wedge (\exists x.\ P2(x)) \Longrightarrow$
$(\forall x.\ P3(x) \longrightarrow P0(x)) \wedge (\exists x.\ P3(x)) \Longrightarrow$
$(\forall x.\ P4(x) \longrightarrow P0(x)) \wedge (\exists x.\ P4(x)) \Longrightarrow$
$(\forall x.\ P5(x) \longrightarrow P0(x)) \wedge (\exists x.\ P5(x)) \Longrightarrow$
$(\exists x.\ Q1(x)) \wedge (\forall x.\ Q1(x) \longrightarrow Q0(x)) \Longrightarrow$
$\forall x.\ P0(x) \longrightarrow ((\forall y.\ Q0(y) \longrightarrow R(x,y)) \vee$
$(\forall y.\ P0(y) \wedge S(y,x) \wedge (\exists z.\ Q0(z) \wedge R(y,z)) \longrightarrow R(x,y))) \Longrightarrow$
$\forall x\ y.\ P3(y) \wedge (P5(x) \vee P4(x)) \longrightarrow S(x,y) \Longrightarrow$
$\forall x\ y.\ P3(x) \wedge P2(y) \longrightarrow S(x,y) \Longrightarrow$
$\forall x\ y.\ P2(x) \wedge P1(y) \longrightarrow S(x,y) \Longrightarrow$
$\forall x\ y.\ P1(x) \wedge (P2(y) \vee Q1(y)) \longrightarrow \neg R(x,y) \Longrightarrow$
$\forall x\ y.\ P3(x) \wedge P4(y) \longrightarrow R(x,y) \Longrightarrow$
$\forall x\ y.\ P3(x) \wedge P5(y) \longrightarrow \neg R(x,y) \Longrightarrow$
$\forall x.\ P4(x) \vee P5(x) \longrightarrow (\exists y.\ Q0(y) \wedge R(x,y)) \Longrightarrow$
$\exists x\ y.\ P0(x) \wedge P0(y) \wedge (\exists z.\ Q1(z) \wedge R(y,z) \wedge R(x,y))$
⟨*proof*⟩

theorem *p48*: $a = b \vee c = d \Longrightarrow a = c \vee b = d \Longrightarrow a = d \vee b = c$ ⟨*proof*⟩

theorem *p49*: $\exists x\ y.\ \forall (z::'a).\ z = x \vee z = y \Longrightarrow P(a) \wedge P(b) \Longrightarrow (a::'a) \neq b \Longrightarrow \forall x.\ P(x)$
⟨*proof*⟩

theorem *p50*: $\forall x.\ F(a,x) \vee (\forall y.\ F(x,y)) \Longrightarrow \exists x.\ \forall y.\ F(x,y)$

⟨*proof*⟩

theorem *p51*: $\exists\, z\ w.\ \forall\, x\ y.\ F(x,y) \longleftrightarrow x = z \wedge y = w \Longrightarrow$
$\exists\, z.\ \forall\, x.\ (\exists\, w.\ \forall\, y.\ F(x,y) \longleftrightarrow y = w) \longleftrightarrow x = z$
⟨*proof*⟩

theorem *p52*: $\exists\, z\ w.\ \forall\, x\ y.\ F(x,y) \longleftrightarrow x = z \wedge y = w \Longrightarrow$
$\exists\, w.\ \forall\, y.\ (\exists\, z.\ \forall\, x.\ F(x,y) \longleftrightarrow x = z) \longleftrightarrow y = w$
⟨*proof*⟩

theorem *p55*:
$\exists\, x.\ L(x) \wedge K(x,a) \Longrightarrow$
$L(a) \wedge L(b) \wedge L(c) \Longrightarrow$
$\forall\, x.\ L(x) \longrightarrow x = a \vee x = b \vee x = c \Longrightarrow$
$\forall\, y\ x.\ K(x,y) \longrightarrow H(x,y) \Longrightarrow$
$\forall\, x\ y.\ K(x,y) \longrightarrow \neg R(x,y) \Longrightarrow$
$\forall\, x.\ H(a,x) \longrightarrow \neg H(c,x) \Longrightarrow$
$\forall\, x.\ x \neq b \longrightarrow H(a,x) \Longrightarrow$
$\forall\, x.\ \neg R(x,a) \longrightarrow H(b,x) \Longrightarrow$
$\forall\, x.\ H(a,x) \longrightarrow H(b,x) \Longrightarrow$ — typo in text
$\forall\, x.\ \exists\, y.\ \neg H(x,y) \Longrightarrow$
$a \neq b \Longrightarrow$
$K(a,a)$
⟨*proof*⟩

theorem *p56*: $(\forall\, x.\ (\exists\, y.\ F(y) \wedge x = f(y)) \longrightarrow F(x)) \longleftrightarrow (\forall\, x.\ F(x) \longrightarrow F(f(x)))$
⟨*proof*⟩

theorem *p57*: $F(f(a,b),f(b,c)) \Longrightarrow F(f(b,c),f(a,c)) \Longrightarrow$
$\forall\, x\ y\ z.\ F(x,y) \wedge F(y,z) \longrightarrow F(x,z) \Longrightarrow F(f(a,b),f(a,c))$ ⟨*proof*⟩

theorem *p58*: $\forall\, x\ y.\ f(x) = g(y) \Longrightarrow \forall\, x\ y.\ f(f(x)) = f(g(y))$
⟨*proof*⟩

theorem *p59*: $\forall\, x{::}'a.\ F(x) \longleftrightarrow \neg F(f(x)) \Longrightarrow \exists\, x.\ F(x) \wedge \neg F(f(x))$
⟨*proof*⟩

theorem *p60*: $\forall\, x.\ F(x,f(x)) \longleftrightarrow (\exists\, y.\ (\forall\, z.\ F(z,y) \longrightarrow F(z,f(x))) \wedge F(x,y))$
⟨*proof*⟩

theorem *p61*: $\forall\, x\ y\ z.\ f(x,f(y,z)) = f(f(x,y),z) \Longrightarrow \forall\, x\ y\ z\ w.\ f(x,f(y,f(z,w))) = f(f(f(x,y),z),w)$
⟨*proof*⟩

end

3 Primes

theory *Primes-Ex*

imports *Auto2-Main*
begin

3.1 Basic definition

definition *prime* :: *nat* ⇒ *bool* **where** [*rewrite*]:
$prime\ p = (1 < p \wedge (\forall m.\ m\ dvd\ p \longrightarrow m = 1 \vee m = p))$

lemma *primeD1* [*forward*]: $prime\ p \Longrightarrow 1 < p$ ⟨*proof*⟩
lemma *primeD2*: $prime\ p \Longrightarrow m\ dvd\ p \Longrightarrow m = 1 \vee m = p$ ⟨*proof*⟩
⟨*ML*⟩

theorem *exists-prime* [*resolve*]: $\exists p.\ prime\ p$
⟨*proof*⟩

lemma *prime-odd-nat*: $prime\ p \Longrightarrow p > 2 \Longrightarrow odd\ p$ ⟨*proof*⟩

lemma *prime-imp-coprime-nat* [*backward2*]: $prime\ p \Longrightarrow \neg\ p\ dvd\ n \Longrightarrow coprime\ p\ n$ ⟨*proof*⟩

lemma *prime-dvd-mult-nat*: $prime\ p \Longrightarrow p\ dvd\ m * n \Longrightarrow p\ dvd\ m \vee p\ dvd\ n$
⟨*proof*⟩
⟨*ML*⟩

theorem *prime-dvd-intro*: $prime\ p \Longrightarrow p * q = m * n \Longrightarrow p\ dvd\ m \vee p\ dvd\ n$
⟨*proof*⟩
⟨*ML*⟩

lemma *prime-dvd-mult-eq-nat*: $prime\ p \Longrightarrow p\ dvd\ m * n = (p\ dvd\ m \vee p\ dvd\ n)$
⟨*proof*⟩

lemma *not-prime-eq-prod-nat* [*backward1*]: $n > 1 \Longrightarrow \neg\ prime\ n \Longrightarrow$
$\exists m\ k.\ n = m * k \wedge 1 < m \wedge m < n \wedge 1 < k \wedge k < n$
⟨*proof*⟩

lemma *prime-dvd-power-nat*: $prime\ p \Longrightarrow p\ dvd\ x\hat{}n \Longrightarrow p\ dvd\ x$ ⟨*proof*⟩
⟨*ML*⟩

lemma *prime-dvd-power-nat-iff*: $prime\ p \Longrightarrow n > 0 \Longrightarrow p\ dvd\ x\hat{}n \longleftrightarrow p\ dvd\ x$
⟨*proof*⟩

lemma *prime-nat-code*: $prime\ p = (1 < p \wedge (\forall x.\ 1 < x \wedge x < p \longrightarrow \neg\ x\ dvd\ p))$
⟨*proof*⟩

lemma *prime-factor-nat* [*backward*]: $n \neq 1 \Longrightarrow \exists p.\ p\ dvd\ n \wedge prime\ p$
⟨*proof*⟩

lemma *prime-divprod-pow-nat*:

$prime\ p \Longrightarrow coprime\ a\ b \Longrightarrow p\hat{}n\ dvd\ a * b \Longrightarrow p\hat{}n\ dvd\ a \vee p\hat{}n\ dvd\ b$ ⟨*proof*⟩

lemma *prime-product* [*forward*]: $prime\ (p * q) \Longrightarrow p = 1 \vee q = 1$
⟨*proof*⟩

lemma *prime-exp*: $prime\ (p\ \hat{}\ n) \longleftrightarrow n = 1 \wedge prime\ p$ ⟨*proof*⟩

lemma *prime-power-mult*: $prime\ p \Longrightarrow x * y = p\ \hat{}\ k \Longrightarrow \exists\, i\ j.\ x = p\ \hat{}\ i \wedge y = p\ \hat{}\ j$
⟨*proof*⟩

3.2 Infinitude of primes

theorem *bigger-prime* [*resolve*]: $\exists\, p.\ prime\ p \wedge n < p$
⟨*proof*⟩

theorem *primes-infinite*: $\neg\ finite\ \{p.\ prime\ p\}$
⟨*proof*⟩

3.3 Existence and uniqueness of prime factorization

theorem *factorization-exists*: $n > 0 \Longrightarrow \exists\, M.\ (\forall\, p{\in}\#M.\ prime\ p) \wedge n = (\prod i{\in}\#M.\ i)$
⟨*proof*⟩

theorem *prime-dvd-multiset* [*backward1*]: $prime\ p \Longrightarrow p\ dvd\ (\prod i{\in}\#M.\ i) \Longrightarrow \exists\, n.\ n{\in}\#M \wedge p\ dvd\ n$
⟨*proof*⟩

theorem *factorization-unique-aux*:
$\forall\, p{\in}\#M.\ prime\ p \Longrightarrow \forall\, p{\in}\#N.\ prime\ p \Longrightarrow (\prod i{\in}\#M.\ i)\ dvd\ (\prod i{\in}\#N.\ i) \Longrightarrow M \subseteq\#\ N$
⟨*proof*⟩
⟨*ML*⟩

theorem *factorization-unique*:
$\forall\, p{\in}\#M.\ prime\ p \Longrightarrow \forall\, p{\in}\#N.\ prime\ p \Longrightarrow (\prod i{\in}\#M.\ i) = (\prod i{\in}\#N.\ i) \Longrightarrow M = N$
⟨*proof*⟩
⟨*ML*⟩

end

References

[1] F. J. Pelletier. Seventy-five problems for testing automatic theorem provers. *Journal of Automated Reasoning*, 2:191–216, 1986.

[2] B. Zhan. Auto2: a saturation-based heuristic prover for higher-order logic. In J. C. Blanchette and S. Merz, editors, *ITP 2016*, pages 441–456, 2016.