A Proof from THE BOOK: The Partial Fraction Expansion of the Cotangent

Manuel Eberl

September 9, 2026

Abstract

In this article, I formalise a proof from THE BOOK [1, Chapter 23]; namely a formula that was called 'one of the most beautiful formulas involving elementary functions':

$$\pi \cot(\pi z) = \frac{1}{z} + \sum_{n=1}^{\infty} \left(\frac{1}{z+n} + \frac{1}{z-n} \right)$$

The proof uses Herglotz's trick to show the real case and analytic continuation for the complex case.

Contents

1 The Partial-Fraction Formula for the Cotangent Function **2**
1.1 Auxiliary lemmas 2
1.2 Definition of auxiliary function 3
1.3 Holomorphicity and continuity 4
1.4 Functional equations 5
1.5 The limit at 0 6
1.6 Final result 6

1 The Partial-Fraction Formula for the Cotangent Function

theory *Cotangent-PFD-Formula*
imports *HOL−Complex-Analysis.Complex-Analysis HOL−Real-Asymp.Real-Asymp*

begin

1.1 Auxiliary lemmas

lemma *uniformly-on-image*:
uniformly-on $(f \mathbin{`} A)$ g = *filtercomap* $(\lambda h.\ h \circ f)$ (*uniformly-on* A $(g \circ f)$)
⟨*proof*⟩

lemma *uniform-limit-image*:
uniform-limit $(f \mathbin{`} A)$ g h $F \longleftrightarrow$ *uniform-limit* A $(\lambda x\ y.\ g\ x\ (f\ y))$ $(\lambda x.\ h\ (f\ x))$ F
⟨*proof*⟩

lemma *Ints-add-iff1* [*simp*]: $x \in \mathbb{Z} \Longrightarrow x + y \in \mathbb{Z} \longleftrightarrow y \in \mathbb{Z}$
⟨*proof*⟩

lemma *Ints-add-iff2* [*simp*]: $y \in \mathbb{Z} \Longrightarrow x + y \in \mathbb{Z} \longleftrightarrow x \in \mathbb{Z}$
⟨*proof*⟩

If a set is discrete (i.e. the difference between any two points is bounded from below), it has no limit points:

lemma *discrete-imp-not-islimpt*:
assumes *e*: $0 < e$
and *d*: $\forall x \in S.\ \forall y \in S.\ \mathit{dist}\ y\ x < e \longrightarrow y = x$
shows $\neg x$ *islimpt* S
⟨*proof*⟩

In particular, the integers have no limit point:

lemma *Ints-not-limpt*: $\neg((x :: 'a :: \mathit{real\text{-}normed\text{-}algebra\text{-}1})\ \mathit{islimpt}\ \mathbb{Z})$
⟨*proof*⟩

The following lemma allows evaluating telescoping sums of the form

$$\sum_{n=0}^{\infty} (f(n) - f(n+k))$$

where $f(n) \longrightarrow 0$, i.e. where all terms except for the first k are cancelled by later summands.

lemma *sums-long-telescope*:
fixes $f :: \mathit{nat} \Rightarrow 'a :: \{\mathit{topological\text{-}group\text{-}add}, \mathit{topological\text{-}comm\text{-}monoid\text{-}add}, \mathit{ab\text{-}group\text{-}add}\}$
assumes *lim*: $f \longrightarrow 0$
shows $(\lambda n.\ f\ n - f\ (n + c))$ *sums* $(\sum k{<}c.\ f\ k)$ (**is** - *sums* *?S*)
⟨*proof*⟩

1.2 Definition of auxiliary function

The following function is the infinite sum appearing on the right-hand side of the cotangent formula. It can be written either as

$$\sum_{n=1}^{\infty}\left(\frac{1}{x+n}+\frac{1}{x-n}\right)$$

or as

$$2x\sum_{n=1}^{\infty}\frac{1}{x^2-n^2}\ .$$

definition *cot-pfd* :: *'a* :: {*real-normed-field*, *banach*} ⇒ *'a* **where**
cot-pfd x = (∑ *n*. *2* ∗ *x* / (*x* ^ *2* − *of-nat* (*Suc n*) ^ *2*))

The sum in the definition of *cot-pfd* converges uniformly on compact sets. This implies, in particular, that *cot-pfd* is holomorphic (and thus also continuous).

lemma *uniform-limit-cot-pfd-complex*:
assumes *R* ≥ *0*
shows *uniform-limit* (*cball 0 R* :: *complex set*)
(λ*N x*. ∑ *n*<*N*. *2* ∗ *x* / (*x* ^ *2* − *of-nat* (*Suc n*) ^ *2*)) *cot-pfd sequentially*
⟨*proof*⟩

lemma *sums-cot-pfd-complex*:
fixes *x* :: *complex*
shows (λ*n*. *2* ∗ *x* / (*x* ^ *2* − *of-nat* (*Suc n*) ^ *2*)) *sums cot-pfd x*
⟨*proof*⟩

lemma *sums-cot-pfd-complex'-aux*:
fixes *x* :: *'a* :: {*banach*, *real-normed-field*, *field-char-0*}
assumes *x* ∉ ℤ − {*0*}
shows *2* ∗ *x* / (*x* ^ *2* − *of-nat* (*Suc n*) ^ *2*) =
1 / (*x* + *of-nat* (*Suc n*)) + *1* / (*x* − *of-nat* (*Suc n*))
⟨*proof*⟩

lemma *sums-cot-pfd-complex'*:
fixes *x* :: *complex*
assumes *x* ∉ ℤ − {*0*}
shows (λ*n*. *1* / (*x* + *of-nat* (*Suc n*)) + *1* / (*x* − *of-nat* (*Suc n*))) *sums cot-pfd x*
⟨*proof*⟩

lemma *summable-cot-pfd-complex*:
fixes *x* :: *complex*
shows *summable* (λ*n*. *2* ∗ *x* / (*x* ^ *2* − *of-nat* (*Suc n*) ^ *2*))
⟨*proof*⟩

lemma *summable-cot-pfd-real*:
fixes x :: *real*
shows *summable* ($\lambda n.\ 2 * x\ /\ (x\ \hat{}\ 2 - of\text{-}nat\ (Suc\ n)\ \hat{}\ 2)$)
⟨*proof*⟩

lemma *sums-cot-pfd-real*:
fixes x :: *real*
shows ($\lambda n.\ 2 * x\ /\ (x\ \hat{}\ 2 - of\text{-}nat\ (Suc\ n)\ \hat{}\ 2)$) *sums cot-pfd x*
⟨*proof*⟩

lemma *cot-pfd-complex-of-real* [*simp*]: *cot-pfd* (*complex-of-real x*) = *of-real* (*cot-pfd x*)
⟨*proof*⟩

lemma *uniform-limit-cot-pfd-real*:
assumes $R \geq 0$
shows *uniform-limit* (*cball 0 R* :: *real set*)
($\lambda N\ x.\ \sum n{<}N.\ 2 * x\ /\ (x\ \hat{}\ 2 - of\text{-}nat\ (Suc\ n)\ \hat{}\ 2)$) *cot-pfd sequentially*
⟨*proof*⟩

1.3 Holomorphicity and continuity

lemma *has-field-derivative-cot-pfd-complex*:
fixes z :: *complex*
assumes z: $z \in -(\mathbb{Z}{-}\{0\})$
shows (*cot-pfd has-field-derivative* (*−Polygamma 1* (*1 + z*) *− Polygamma 1* (*1 − z*))) (*at z*)
⟨*proof*⟩

lemma *has-field-derivative-cot-pfd-complex′* [*derivative-intros*]:
assumes (*g has-field-derivative g′*) (*at x within A*) **and** $g\ x \notin \mathbb{Z} - \{0\}$
shows ((λ*x*. *cot-pfd* (*g x* :: *complex*)) *has-field-derivative*
(*−Polygamma 1* (*1 + g x*) *− Polygamma 1* (*1 − g x*)) * *g′*) (*at x within A*)
⟨*proof*⟩

lemma *Polygamma-real-conv-complex*: $x \neq 0 \Longrightarrow$ *Polygamma n x* = *Re* (*Polygamma n* (*of-real x*))
⟨*proof*⟩

lemma *has-field-derivative-cot-pfd-real* [*derivative-intros*]:
assumes (*g has-field-derivative g′*) (*at x within A*) **and** $g\ x \notin \mathbb{Z} - \{0\}$
shows ((λ*x*. *cot-pfd* (*g x* :: *real*)) *has-field-derivative*
(*−Polygamma 1* (*1 + g x*) *− Polygamma 1* (*1 − g x*)) * *g′*) (*at x within A*)
⟨*proof*⟩

lemma *holomorphic-on-cot-pfd* [*holomorphic-intros*]:
assumes $A \subseteq -(\mathbb{Z}{-}\{0\})$

shows *cot-pfd holomorphic-on A*
⟨*proof*⟩

lemma *holomorphic-on-cot-pfd'* [*holomorphic-intros*]:
assumes *f holomorphic-on A* $\bigwedge x.\ x \in A \Longrightarrow f\ x \notin \mathbb{Z} - \{0\}$
shows $(\lambda x.\ \textit{cot-pfd}\ (f\ x))$ *holomorphic-on A*
⟨*proof*⟩

lemma *continuous-on-cot-pfd-complex* [*continuous-intros*]:
assumes *continuous-on A f* $\bigwedge z.\ z \in A \Longrightarrow f\ z \notin \mathbb{Z} - \{0\}$
shows *continuous-on A* $(\lambda x.\ \textit{cot-pfd}\ (f\ x :: \textit{complex}))$
⟨*proof*⟩

lemma *continuous-on-cot-pfd-real* [*continuous-intros*]:
assumes *continuous-on A f* $\bigwedge z.\ z \in A \Longrightarrow f\ z \notin \mathbb{Z} - \{0\}$
shows *continuous-on A* $(\lambda x.\ \textit{cot-pfd}\ (f\ x :: \textit{real}))$
⟨*proof*⟩

1.4 Functional equations

In this section, we will show three few functional equations for the function *cot-pfd*. The first one is trivial; the other two are a bit tedious and not very insightful, so I will not comment on them.

cot-pfd is an odd function:

lemma *cot-pfd-complex-minus* [*simp*]: *cot-pfd* $(-x :: \textit{complex}) = -\textit{cot-pfd}\ x$
⟨*proof*⟩

lemma *cot-pfd-real-minus* [*simp*]: *cot-pfd* $(-x :: \textit{real}) = -\textit{cot-pfd}\ x$
⟨*proof*⟩

1 / x + cot-pfd x is periodic with period 1:

lemma *cot-pfd-plus-1-complex*:
assumes $x \notin \mathbb{Z}$
shows *cot-pfd* $(x + 1 :: \textit{complex}) = \textit{cot-pfd}\ x - 1 / (x + 1) + 1 / x$
⟨*proof*⟩

lemma *cot-pfd-plus-1-real*:
assumes $x \notin \mathbb{Z}$
shows *cot-pfd* $(x + 1 :: \textit{real}) = \textit{cot-pfd}\ x - 1 / (x + 1) + 1 / x$
⟨*proof*⟩

cot-pfd satisfies the following functional equation:

$$2f(x) = f\left(\frac{x}{2}\right) + f\left(\frac{x+1}{2}\right) + \frac{2}{x+1}$$

lemma *cot-pfd-funeq-complex*:

fixes x :: *complex*
assumes $x \notin \mathbb{Z}$
shows $2 * \textit{cot-pfd}\ x = \textit{cot-pfd}\ (x\ /\ 2) + \textit{cot-pfd}\ ((x + 1)\ /\ 2) + 2\ /\ (x + 1)$
⟨*proof*⟩

lemma *cot-pfd-funeq-real*:
fixes x :: *real*
assumes $x \notin \mathbb{Z}$
shows $2 * \textit{cot-pfd}\ x = \textit{cot-pfd}\ (x\ /\ 2) + \textit{cot-pfd}\ ((x + 1)\ /\ 2) + 2\ /\ (x + 1)$
⟨*proof*⟩

1.5 The limit at 0

lemma *cot-pfd-real-tendsto-0*: $\textit{cot-pfd} \longrightarrow 0 \rightarrow (0 :: \textit{real})$
⟨*proof*⟩

1.6 Final result

To show the final result, we first prove the real case using Herglotz's trick, following the presentation in 'Proofs from THE BOOK'. [1, Chapter 23].

lemma *cot-pfd-formula-real*:
assumes $x \notin \mathbb{Z}$
shows $pi * \textit{cot}\ (pi * x) = 1\ /\ x + \textit{cot-pfd}\ x$
⟨*proof*⟩

We now lift the result from the domain $\mathbb{R}\backslash\mathbb{Z}$ to $\mathbb{C}\backslash\mathbb{Z}$. We do this by noting that $\mathbb{C}\backslash\mathbb{Z}$ is connected and the point $\frac{1}{2}$ is both in $\mathbb{C}\backslash\mathbb{Z}$ and a limit point of $\mathbb{R}\backslash\mathbb{Z}$.

lemma *one-half-limit-point-Reals-minus-Ints*: $(1\ /\ 2 :: \textit{complex})\ \textit{islimpt}\ \mathbb{R} - \mathbb{Z}$
⟨*proof*⟩

theorem *cot-pfd-formula-complex*:
fixes z :: *complex*
assumes $z \notin \mathbb{Z}$
shows $pi * \textit{cot}\ (pi * z) = 1\ /\ z + \textit{cot-pfd}\ z$
⟨*proof*⟩

end

References

[1] M. Aigner and G. M. Ziegler. *Proofs from THE BOOK*. Springer, 4th edition, 2009.