Laplace Transform

Fabian Immler

April 18, 2024

Abstract

This entry formalizes the Laplace transform and concrete Laplace transforms for arithmetic functions, frequency shift, integration and (higher) differentiation in the time domain. It proves Lerch's lemma and uniqueness of the Laplace transform for continuous functions. In order to formalize the foundational assumptions, this entry contains a formalization of piecewise continuous functions and functions of exponential order.

Contents

1 References **2**

2 Library Additions **2**
2.1 Derivatives 2
2.2 Integrals 2
2.3 Miscellaneous 3

3 Piecewise Continous Functions **4**
3.1 at within filters 4
3.2 intervals 4

4 Existence **9**
4.1 Definition 9
4.2 Condition for Existence: Exponential Order 9
4.3 Concrete Laplace Transforms 11
4.4 higher derivatives 14

5 Lerch Lemma **15**

6 Uniqueness of Laplace Transform **16**

theory *Laplace-Transform-Library*
imports
HOL−Analysis.Analysis
begin

1 References

Much of this formalization is based on Schiff's textbook [3]. Parts of this formalization are inspired by the HOL-Light formalization ([4], [1], [2]), but stated more generally for piecewise continuous (instead of piecewise continuously differentiable) functions.

2 Library Additions

2.1 Derivatives

lemma *DERIV-compose-FDERIV*:— TODO: generalize and move from HOL-ODE
  assumes $DERIV\ f\ (g\ x) :> f'$
  assumes $(g\ \textit{has-derivative}\ g')\ (\textit{at}\ x\ \textit{within}\ s)$
  shows $((\lambda x.\ f\ (g\ x))\ \textit{has-derivative}\ (\lambda x.\ g'\ x * f'))\ (\textit{at}\ x\ \textit{within}\ s)$
  ⟨*proof*⟩

lemmas *has-derivative-sin*[*derivative-intros*] = *DERIV-sin*[*THEN DERIV-compose-FDERIV*]
 and *has-derivative-cos*[*derivative-intros*] = *DERIV-cos*[*THEN DERIV-compose-FDERIV*]
 and *has-derivative-exp*[*derivative-intros*] = *DERIV-exp*[*THEN DERIV-compose-FDERIV*]

2.2 Integrals

lemma *negligible-real-ivlI*:
  fixes $a\ b{::}real$
  assumes $a \geq b$
  shows $negligible\ \{a\ ..\ b\}$
⟨*proof*⟩

lemma *absolutely-integrable-on-combine*:
  fixes $f :: real \Rightarrow 'a{::}euclidean\text{-}space$
  assumes $f\ \textit{absolutely-integrable-on}\ \{a..c\}$
    and $f\ \textit{absolutely-integrable-on}\ \{c..b\}$
    and $a \leq c$
    and $c \leq b$
  shows $f\ \textit{absolutely-integrable-on}\ \{a..b\}$
  ⟨*proof*⟩

lemma *dominated-convergence-at-top*:
  fixes $f :: real \Rightarrow 'n{::}euclidean\text{-}space \Rightarrow 'm{::}euclidean\text{-}space$
  assumes f: $\bigwedge k.\ (f\ k)\ \textit{integrable-on}\ s$ **and** h: $h\ \textit{integrable-on}\ s$
    and le: $\bigwedge k\ x.\ x \in s \Longrightarrow norm\ (f\ k\ x) \leq h\ x$
    and $conv$: $\forall x \in s.\ ((\lambda k.\ f\ k\ x) \longrightarrow g\ x)\ \textit{at-top}$
  shows $g\ \textit{integrable-on}\ s\ ((\lambda k.\ integral\ s\ (f\ k)) \longrightarrow integral\ s\ g)\ \textit{at-top}$
⟨*proof*⟩

lemma *has-integral-dominated-convergence-at-top*:

fixes f :: *real* $\Rightarrow$ $'n$::*euclidean-space* $\Rightarrow$ $'m$::*euclidean-space*
assumes $\bigwedge k.$ (f k *has-integral* y k) s h *integrable-on* s
$\bigwedge k\ x.\ x \in s \Longrightarrow$ *norm* (f k x) $\leq h\ x\ \forall x \in s.$ (($\lambda k.\ f\ k\ x$) $\longrightarrow g\ x$) *at-top*
and x: ($y \longrightarrow x$) *at-top*
shows (g *has-integral* x) s
⟨*proof*⟩

lemma *integral-indicator-eq-restriction*:
fixes f::$'a$::*euclidean-space* $\Rightarrow$ $'b$::*banach*
assumes f: f *integrable-on* R
and $R \subseteq S$
shows *integral* S ($\lambda x.$ *indicator* R $x *_R f\ x$) = *integral* R f
⟨*proof*⟩

lemma
improper-integral-at-top:
fixes f::*real* $\Rightarrow$ $'a$::*euclidean-space*
assumes f *absolutely-integrable-on* $\{a..\}$
shows (($\lambda x.$ *integral* $\{a..x\}$ f) $\longrightarrow$ *integral* $\{a..\}$ f) *at-top*
⟨*proof*⟩

lemma *norm-integrable-onI*: ($\lambda x.$ *norm* ($f\ x$)) *integrable-on* S
if f *absolutely-integrable-on* S
for f::$'a$::*euclidean-space*$\Rightarrow$$'b$::*euclidean-space*
⟨*proof*⟩

lemma
has-integral-improper-at-topI:
fixes f::*real* $\Rightarrow$ $'a$::*banach*
assumes I: $\forall_F$ k *in at-top.* (f *has-integral* I k) $\{a..k\}$
assumes J: ($I \longrightarrow J$) *at-top*
shows (f *has-integral* J) $\{a..\}$
⟨*proof*⟩

lemma *has-integral-improperE*:
fixes f::*real* $\Rightarrow$ $'a$::*euclidean-space*
assumes I: (f *has-integral* I) $\{a..\}$
assumes *ai*: f *absolutely-integrable-on* $\{a..\}$
obtains J **where**
$\bigwedge k.$ (f *has-integral* J k) $\{a..k\}$
($J \longrightarrow I$) *at-top*
⟨*proof*⟩

2.3 Miscellaneous

lemma *AE-BallI*: *AE* $x \in X$ *in* F. $P\ x$ **if** $\forall x \in X.\ P\ x$
⟨*proof*⟩

lemma *bounded-le-Sup*:

assumes *bounded* $(f\ `\ S)$
shows $\forall x{\in}S.\ norm\ (f\ x) \leq Sup\ (norm\ `\ f\ `\ S)$
⟨*proof*⟩

end

3 Piecewise Continous Functions

theory *Piecewise-Continuous*
imports
Laplace-Transform-Library
begin

3.1 at within filters

lemma *at-within-self-singleton*[*simp*]: *at i within* $\{i\} = bot$
⟨*proof*⟩

lemma *at-within-t1-space-avoid*:
$(at\ x\ within\ X - \{i\}) = (at\ x\ within\ X)$ **if** $x \neq i$ **for** $x\ i{::}'a{::}t1\text{-}space$
⟨*proof*⟩

lemma *at-within-t1-space-avoid-finite*:
$(at\ x\ within\ X - I) = (at\ x\ within\ X)$ **if** $finite\ I\ x \notin I$ **for** $x{::}'a{::}t1\text{-}space$
⟨*proof*⟩

lemma *at-within-interior*:
$NO\text{-}MATCH\ (UNIV{::}'a\ set)\ (S{::}'a{::}topological\text{-}space\ set) \Longrightarrow x \in interior\ S \Longrightarrow at\ x\ within\ S = at\ x$
⟨*proof*⟩

3.2 intervals

lemma *Compl-Icc*: $-\ \{a\ ..\ b\} = \{..{<}a\} \cup \{b{<}..\}$ **for** $a\ b{::}'a{::}linorder$
⟨*proof*⟩

lemma *interior-Icc*[*simp*]: $interior\ \{a..b\} = \{a{<}..{<}b\}$
for $a\ b{::}'a{::}\{linorder\text{-}topology,\ dense\text{-}order,\ no\text{-}bot,\ no\text{-}top\}$
— TODO: is *no-bot* and *no-top* really required?
⟨*proof*⟩

lemma *closure-finite*[*simp*]: $closure\ X = X$ **if** $finite\ X$ **for** $X{::}'a{::}t1\text{-}space\ set$
⟨*proof*⟩

definition *piecewise-continuous-on* $::\ 'a{::}linorder\text{-}topology \Rightarrow 'a \Rightarrow 'a\ set \Rightarrow ('a \Rightarrow 'b{::}topological\text{-}space) \Rightarrow bool$
where $piecewise\text{-}continuous\text{-}on\ a\ b\ I\ f \longleftrightarrow$
$(continuous\text{-}on\ (\{a\ ..\ b\} - I)\ f \wedge finite\ I \wedge$
$(\forall i{\in}I.\ (i \in \{a{<}..b\} \longrightarrow (\exists l.\ (f \longrightarrow l)\ (at\text{-}left\ i))) \wedge$

$(i \in \{a..<b\} \longrightarrow (\exists\, u.\ (f \longrightarrow u)\ (\textit{at-right}\ i)))))$

lemma *piecewise-continuous-on-subset*:
piecewise-continuous-on a b I f $\Longrightarrow \{c\ ..\ d\} \subseteq \{a\ ..\ b\} \Longrightarrow$ *piecewise-continuous-on c d I f*
⟨*proof*⟩

lemma *piecewise-continuous-onE*:
assumes *piecewise-continuous-on a b I f*
obtains *l u*
where *finite I*
and *continuous-on* $(\{a..b\} - I)\ f$
and $(\bigwedge i.\ i \in I \Longrightarrow a < i \Longrightarrow i \leq b \Longrightarrow (f \longrightarrow l\ i)\ (\textit{at-left}\ i))$
and $(\bigwedge i.\ i \in I \Longrightarrow a \leq i \Longrightarrow i < b \Longrightarrow (f \longrightarrow u\ i)\ (\textit{at-right}\ i))$
⟨*proof*⟩

lemma *piecewise-continuous-onI*:
assumes *finite I continuous-on* $(\{a..b\} - I)\ f$
and $(\bigwedge i.\ i \in I \Longrightarrow a < i \Longrightarrow i \leq b \Longrightarrow (f \longrightarrow l\ i)\ (\textit{at-left}\ i))$
and $(\bigwedge i.\ i \in I \Longrightarrow a \leq i \Longrightarrow i < b \Longrightarrow (f \longrightarrow u\ i)\ (\textit{at-right}\ i))$
shows *piecewise-continuous-on a b I f*
⟨*proof*⟩

lemma *piecewise-continuous-onI′*:
fixes *a b*::*′a*::{*linorder-topology*, *dense-order*, *no-bot*, *no-top*}
assumes *finite I* $\bigwedge x.\ a < x \Longrightarrow x < b \Longrightarrow$ *isCont f x*
and $a \notin I \Longrightarrow$ *continuous* (*at-right a*) *f*
and $b \notin I \Longrightarrow$ *continuous* (*at-left b*) *f*
and $(\bigwedge i.\ i \in I \Longrightarrow a < i \Longrightarrow i \leq b \Longrightarrow (f \longrightarrow l\ i)\ (\textit{at-left}\ i))$
and $(\bigwedge i.\ i \in I \Longrightarrow a \leq i \Longrightarrow i < b \Longrightarrow (f \longrightarrow u\ i)\ (\textit{at-right}\ i))$
shows *piecewise-continuous-on a b I f*
⟨*proof*⟩

lemma *piecewise-continuous-onE′*:
fixes *a b*::*′a*::{*linorder-topology*, *dense-order*, *no-bot*, *no-top*}
assumes *piecewise-continuous-on a b I f*
obtains *l u*
where *finite I*
and $\bigwedge x.\ a < x \Longrightarrow x < b \Longrightarrow x \notin I \Longrightarrow$ *isCont f x*
and $(\bigwedge x.\ a < x \Longrightarrow x \leq b \Longrightarrow (f \longrightarrow l\ x)\ (\textit{at-left}\ x))$
and $(\bigwedge x.\ a \leq x \Longrightarrow x < b \Longrightarrow (f \longrightarrow u\ x)\ (\textit{at-right}\ x))$
and $\bigwedge x.\ a \leq x \Longrightarrow x \leq b \Longrightarrow x \notin I \Longrightarrow f\ x = l\ x$
and $\bigwedge x.\ a \leq x \Longrightarrow x \leq b \Longrightarrow x \notin I \Longrightarrow f\ x = u\ x$
⟨*proof*⟩

lemma *tendsto-avoid-at-within*:
$(f \longrightarrow l)$ (*at x within X*)
if $(f \longrightarrow l)$ (*at x within* $X - \{x\}$)
⟨*proof*⟩

lemma *tendsto-within-subset-eventuallyI*:
$(f \longrightarrow fx)$ $(at\ x\ within\ X)$
if g: $(g \longrightarrow gy)$ $(at\ y\ within\ Y)$
and ev: $\forall_F\ x\ in\ (at\ y\ within\ Y).\ f\ x = g\ x$
and xy: $x = y$
and $fxgy$: $fx = gy$
and XY: $X - \{x\} \subseteq Y$
⟨*proof*⟩

lemma *piecewise-continuous-on-insertE*:
assumes *piecewise-continuous-on a b* (*insert i I*) *f*
assumes $i \in \{a\ ..\ b\}$
obtains *g h* **where**
piecewise-continuous-on a i I g
piecewise-continuous-on i b I h
$\bigwedge x.\ a \le x \Longrightarrow x < i \Longrightarrow g\ x = f\ x$
$\bigwedge x.\ i < x \Longrightarrow x \le b \Longrightarrow h\ x = f\ x$
⟨*proof*⟩

lemma *eventually-avoid-finite*:
$\forall_F\ x\ in\ at\ y\ within\ Y.\ x \notin I$ **if** *finite I* **for** *y::'a::t1-space*
⟨*proof*⟩

lemma *eventually-at-left-linorder*:— TODO: generalize $?b < ?a \Longrightarrow \forall_F\ x\ in\ at\text{-}left\ ?a.\ x \in \{?b<..<?a\}$
$a > (b :: 'a :: linorder\text{-}topology) \Longrightarrow eventually\ (\lambda x.\ x \in \{b<..<a\})\ (at\text{-}left\ a)$
⟨*proof*⟩

lemma *eventually-at-right-linorder*:— TODO: generalize $?a < ?b \Longrightarrow \forall_F\ x\ in\ at\text{-}right\ ?a.\ x \in \{?a<..<?b\}$
$a > (b :: 'a :: linorder\text{-}topology) \Longrightarrow eventually\ (\lambda x.\ x \in \{b<..<a\})\ (at\text{-}right\ b)$
⟨*proof*⟩

lemma *piecewise-continuous-on-congI*:
piecewise-continuous-on a b I g
if *piecewise-continuous-on a b I f*
and *eq*: $\bigwedge x.\ x \in \{a\ ..\ b\} - I \Longrightarrow g\ x = f\ x$
⟨*proof*⟩

lemma *piecewise-continuous-on-cong*[*cong*]:
piecewise-continuous-on a b I f $\longleftrightarrow$ *piecewise-continuous-on c d J g*
if $a = c$
$b = d$
$I = J$
$\bigwedge x.\ c \le x \Longrightarrow x \le d \Longrightarrow x \notin J \Longrightarrow f\ x = g\ x$
⟨*proof*⟩

lemma *tendsto-at-left-continuous-on-avoidI*: $(f \longrightarrow g\ i)$ $(at\text{-}left\ i)$

if g: *continuous-on* $(\{a..i\} - I)$ g
and gf: $\bigwedge x.\ a < x \Longrightarrow x < i \Longrightarrow g\ x = f\ x$
$i \notin I$ *finite* I $a < i$
for i::$'a$::*linorder-topology*
⟨*proof*⟩

lemma *tendsto-at-right-continuous-on-avoidI*: $(f \longrightarrow g\ i)$ (*at-right* i)
if g: *continuous-on* $(\{i..b\} - I)$ g
and gf: $\bigwedge x.\ i < x \Longrightarrow x < b \Longrightarrow g\ x = f\ x$
$i \notin I$ *finite* I $i < b$
for i::$'a$::*linorder-topology*
⟨*proof*⟩

lemma *piecewise-continuous-on-insert-leftI*:
piecewise-continuous-on a b (*insert* a I) f **if** *piecewise-continuous-on* a b I f
⟨*proof*⟩

lemma *piecewise-continuous-on-insert-rightI*:
piecewise-continuous-on a b (*insert* b I) f **if** *piecewise-continuous-on* a b I f
⟨*proof*⟩

theorem *piecewise-continuous-on-induct*[*consumes 1*, *case-names empty combine weaken*]:
assumes *pc*: *piecewise-continuous-on* a b I f
assumes *1*: $\bigwedge a\ b\ f.$ *continuous-on* $\{a\ ..\ b\}$ $f \Longrightarrow P\ a\ b\ \{\}\ f$
assumes *2*: $\bigwedge a\ i\ b\ I\ f1\ f2\ f.\ a \le i \Longrightarrow i \le b \Longrightarrow i \notin I \Longrightarrow P\ a\ i\ I\ f1 \Longrightarrow P\ i\ b\ I\ f2 \Longrightarrow$
piecewise-continuous-on a i I $f1 \Longrightarrow$
piecewise-continuous-on i b I $f2 \Longrightarrow$
$(\bigwedge x.\ a \le x \Longrightarrow x < i \Longrightarrow f1\ x = f\ x) \Longrightarrow$
$(\bigwedge x.\ i < x \Longrightarrow x \le b \Longrightarrow f2\ x = f\ x) \Longrightarrow$
$(i > a \Longrightarrow (f \longrightarrow f1\ i)$ (*at-left* i)$) \Longrightarrow$
$(i < b \Longrightarrow (f \longrightarrow f2\ i)$ (*at-right* i)$) \Longrightarrow$
$P\ a\ b$ (*insert* i I) f
assumes *3*: $\bigwedge a\ b\ i\ I\ f.\ P\ a\ b\ I\ f \Longrightarrow$ *finite* $I \Longrightarrow i \notin I \Longrightarrow P\ a\ b$ (*insert* i I) f
shows $P\ a\ b\ I\ f$
⟨*proof*⟩

lemma *continuous-on-imp-piecewise-continuous-on*:
continuous-on $\{a\ ..\ b\}$ $f \Longrightarrow$ *piecewise-continuous-on* a b $\{\}$ f
⟨*proof*⟩

lemma *piecewise-continuous-on-imp-absolutely-integrable*:
fixes a b::*real* **and** f::*real* $\Rightarrow$ $'a$::*euclidean-space*
assumes *piecewise-continuous-on* a b I f
shows f *absolutely-integrable-on* $\{a..b\}$
⟨*proof*⟩

lemma *piecewise-continuous-on-integrable*:

fixes $a\ b$::*real* **and** f::*real* $\Rightarrow$ $'a$::*euclidean-space*
assumes *piecewise-continuous-on* $a\ b\ I\ f$
shows f *integrable-on* $\{a..b\}$
⟨*proof*⟩

lemma *piecewise-continuous-on-comp*:
assumes p: *piecewise-continuous-on* $a\ b\ I\ f$
assumes c: $\bigwedge x.$ *isCont* $(\lambda(x,\ y).\ g\ x\ y)\ x$
shows *piecewise-continuous-on* $a\ b\ I\ (\lambda x.\ g\ x\ (f\ x))$
⟨*proof*⟩

lemma *bounded-piecewise-continuous-image*:
bounded $(f\ `\ \{a\ ..\ b\})$
if *piecewise-continuous-on* $a\ b\ I\ f$ **for** $a\ b$::*real*
⟨*proof*⟩

lemma *tendsto-within-eventually*:
$(f \longrightarrow l)$ (*at* x *within* X)
if
$(f \longrightarrow l)$ (*at* x *within* Y)
$\forall_F\ y$ *in at* x *within* $X.\ y \in Y$
⟨*proof*⟩

lemma *at-within-eq-bot-lemma*:
at x *within* $\{b..c\} = (\textit{if}\ x < b \vee b > c\ \textit{then bot else at}\ x\ \textit{within}\ \{b..c\})$
for $x\ b\ c$::$'a$::*linorder-topology*
⟨*proof*⟩

lemma *at-within-eq-bot-lemma2*:
at x *within* $\{a..b\} = (\textit{if}\ x > b \vee a > b\ \textit{then bot else at}\ x\ \textit{within}\ \{a..b\})$
for $x\ a\ b$::$'a$::*linorder-topology*
⟨*proof*⟩

lemma *piecewise-continuous-on-combine*:
piecewise-continuous-on $a\ c\ J\ f$
if *piecewise-continuous-on* $a\ b\ J\ f$ *piecewise-continuous-on* $b\ c\ J\ f$
⟨*proof*⟩

lemma *piecewise-continuous-on-finite-superset*:
piecewise-continuous-on $a\ b\ I\ f \Longrightarrow I \subseteq J \Longrightarrow$ *finite* $J \Longrightarrow$ *piecewise-continuous-on* $a\ b\ J\ f$
for $a\ b$::$'a$::{*linorder-topology*, *dense-order*, *no-bot*, *no-top*}
⟨*proof*⟩

lemma *piecewise-continuous-on-splitI*:
piecewise-continuous-on $a\ c\ K\ f$
if
piecewise-continuous-on $a\ b\ I\ f$
piecewise-continuous-on $b\ c\ J\ f$

$I \subseteq K$ $J \subseteq K$ *finite* K
for a b::$'a$::{*linorder-topology*, *dense-order*, *no-bot*, *no-top*}
⟨*proof*⟩

end

4 Existence

theory *Existence* **imports**
Piecewise-Continuous
begin

4.1 Definition

definition *has-laplace* :: (*real* ⇒ *complex*) ⇒ *complex* ⇒ *complex* ⇒ *bool*
(**infixr** *has′-laplace 46*)
where (f *has-laplace* L) s ⟷ (($\lambda t.$ *exp* ($t *_R - s$) $*$ f t) *has-integral* L) {*0*..}

lemma *has-laplaceI*:
assumes (($\lambda t.$ *exp* ($t *_R - s$) $*$ f t) *has-integral* L) {*0*..}
shows (f *has-laplace* L) s
⟨*proof*⟩

lemma *has-laplaceD*:
assumes (f *has-laplace* L) s
shows (($\lambda t.$ *exp* ($t *_R - s$) $*$ f t) *has-integral* L) {*0*..}
⟨*proof*⟩

lemma *has-laplace-unique*:
$L = M$ **if**
(f *has-laplace* L) s
(f *has-laplace* M) s
⟨*proof*⟩

4.2 Condition for Existence: Exponential Order

definition *exponential-order* M c f ⟷ $0 < M \wedge (\forall_F$ t *in at-top. norm* (f t) $\leq$ $M * exp$ ($c * t$))

lemma *exponential-orderI*:
assumes $0 < M$ **and** *eo*: $\forall_F$ t *in at-top. norm* (f t) $\leq$ $M * exp$ ($c * t$)
shows *exponential-order* M c f
⟨*proof*⟩

lemma *exponential-orderD*:
assumes *exponential-order* M c f
shows $0 < M$ $\forall_F$ t *in at-top. norm* (f t) $\leq$ $M * exp$ ($c * t$)
⟨*proof*⟩

context
  fixes f::*real* $\Rightarrow$ *complex*
begin

definition *laplace-integrand*::*complex* $\Rightarrow$ *real* $\Rightarrow$ *complex*
  where *laplace-integrand s t* = *exp* $(t *_R - s) * f\ t$

lemma *laplace-integrand-absolutely-integrable-on-Icc*:
  laplace-integrand s absolutely-integrable-on $\{a..b\}$
  if *AE* $x \in \{a..b\}$ *in lebesgue. cmod* $(f\ x) \leq B$ *f integrable-on* $\{a..b\}$
  ⟨*proof*⟩

lemma *laplace-integrand-integrable-on-Icc*:
  laplace-integrand s integrable-on $\{a..b\}$
  if *AE* $x \in \{a..b\}$ *in lebesgue. cmod* $(f\ x) \leq B$ *f integrable-on* $\{a..b\}$
  ⟨*proof*⟩

lemma *eventually-laplace-integrand-le*:
  $\forall_F$ *t in at-top. cmod* (*laplace-integrand s t*) $\leq M * exp\ (-\ (Re\ s - c) * t)$
  if *exponential-order M c f*
  ⟨*proof*⟩

lemma
  assumes *eo*: *exponential-order M c f*
    and *cs*: $c < Re\ s$
  shows *laplace-integrand-integrable-on-Ici-iff*:
    laplace-integrand s integrable-on $\{a..\}$ $\longleftrightarrow$
      ($\forall k > a$. *laplace-integrand s integrable-on* $\{a..k\}$)
    (**is** *?th1*)
  and *laplace-integrand-absolutely-integrable-on-Ici-iff*:
    laplace-integrand s absolutely-integrable-on $\{a..\}$ $\longleftrightarrow$
      ($\forall k > a$. *laplace-integrand s absolutely-integrable-on* $\{a..k\}$)
    (**is** *?th2*)
⟨*proof*⟩

theorem *laplace-exists-laplace-integrandI*:
  assumes *laplace-integrand s integrable-on* $\{0..\}$
  obtains *F* **where** (*f has-laplace F*) *s*
⟨*proof*⟩

lemma
  assumes *eo*: *exponential-order M c f*
    and *pc*: $\bigwedge k$. *AE* $x \in \{0..k\}$ *in lebesgue. cmod* $(f\ x) \leq B\ k$ $\bigwedge k$. *f integrable-on* $\{0..k\}$
    and *s*: $Re\ s > c$
  shows *laplace-integrand-integrable*: *laplace-integrand s integrable-on* $\{0..\}$ (**is** *?th1*)
    and *laplace-integrand-absolutely-integrable*:
      laplace-integrand s absolutely-integrable-on $\{0..\}$ (**is** *?th2*)

⟨*proof*⟩

lemma *piecewise-continuous-on-AE-boundedE*:
assumes *pc*: $\bigwedge k.$ *piecewise-continuous-on a k* (*I k*) *f*
obtains *B* **where** $\bigwedge k.$ *AE* $x \in \{a..k\}$ *in lebesgue. cmod* (*f x*) $\leq$ *B k*
⟨*proof*⟩

theorem *piecewise-continuous-on-has-laplace*:
assumes *eo*: *exponential-order M c f*
and *pc*: $\bigwedge k.$ *piecewise-continuous-on 0 k* (*I k*) *f*
and *s*: *Re s* > *c*
obtains *F* **where** (*f has-laplace F*) *s*
⟨*proof*⟩

end

4.3 Concrete Laplace Transforms

lemma *exp-scaleR-has-vector-derivative-left'*[*derivative-intros*]:
($\lambda t.$ *exp* ($t *_R A$)) *has-vector-derivative* $A *$ *exp* ($t *_R A$)) (*at t within S*)
⟨*proof*⟩

lemma
fixes *a*::*complex*— TODO: generalize
assumes *a*: *0* < *Re a*
shows *integrable-on-cexp-minus-to-infinity*: ($\lambda x.$ *exp* ($x *_R - a$)) *integrable-on* $\{c..\}$
and *integral-cexp-minus-to-infinity*: *integral* $\{c..\}$ ($\lambda x.$ *exp* ($x *_R - a$)) = *exp* ($c *_R - a$) / *a*
⟨*proof*⟩

lemma *has-integral-cexp-minus-to-infinity*:
fixes *a*::*complex*— TODO: generalize
assumes *a*: *0* < *Re a*
shows (($\lambda x.$ *exp* ($x *_R - a$)) *has-integral exp* ($c *_R - a$) / *a*) $\{c..\}$
⟨*proof*⟩

lemma *has-laplace-one*:
((λ-. *1*) *has-laplace inverse s*) *s* **if** *Re s* > *0*
⟨*proof*⟩

lemma *has-laplace-add*:
assumes *f*: (*f has-laplace F*) *S*
assumes *g*: (*g has-laplace G*) *S*
shows (($\lambda x.$ *f x* + *g x*) *has-laplace F* + *G*) *S*
⟨*proof*⟩

lemma *has-laplace-cmul*:
assumes (*f has-laplace F*) *S*

shows $((\lambda x.\ r *_R f\ x)$ *has-laplace* $r *_R F)\ S$
⟨*proof*⟩

lemma *has-laplace-uminus*:
assumes $(f$ *has-laplace* $F)\ S$
shows $((\lambda x.\ -f\ x)$ *has-laplace* $-F)\ S$
⟨*proof*⟩

lemma *has-laplace-minus*:
assumes f: $(f$ *has-laplace* $F)\ S$
assumes g: $(g$ *has-laplace* $G)\ S$
shows $((\lambda x.\ f\ x - g\ x)$ *has-laplace* $F - G)\ S$
⟨*proof*⟩

lemma *has-laplace-spike*:
$(f$ *has-laplace* $L)\ s$
if L: $(g$ *has-laplace* $L)\ s$
and *negligible* T
and $\bigwedge t.\ t \notin T \Longrightarrow t \geq 0 \Longrightarrow f\ t = g\ t$
⟨*proof*⟩

lemma *has-laplace-frequency-shift*:— First Translation Theorem in Schiff
$((\lambda t.\ exp\ (t *_R b) * f\ t)$ *has-laplace* $L)\ s$
if $(f$ *has-laplace* $L)\ (s - b)$
⟨*proof*⟩

theorem *has-laplace-derivative-time-domain*:
$(f'$ *has-laplace* $s * L - f0)\ s$
if L: $(f$ *has-laplace* $L)\ s$
and f': $\bigwedge t.\ t > 0 \Longrightarrow (f$ *has-vector-derivative* $f'\ t)\ (at\ t)$
and $f0$: $(f \longrightarrow f0)\ (at\text{-}right\ 0)$
and eo: *exponential-order* $M\ c\ f$
and cs: $c < Re\ s$
— Proof and statement follow "The Laplace Transform: Theory and Applications" by Joel L. Schiff.
⟨*proof*⟩

lemma *exp-times-has-integral*:
$((\lambda t.\ exp\ (c * t))$ *has-integral* $(if\ c = 0\ then\ t\ else\ exp\ (c * t)\ /\ c) - (if\ c = 0\ then\ t0\ else\ exp\ (c * t0)\ /\ c))\ \{t0\ ..\ t\}$
if $t0 \leq t$
for $c\ t$::*real*
⟨*proof*⟩

lemma *integral-exp-times*:
$integral\ \{t0\ ..\ t\}\ (\lambda t.\ exp\ (c * t)) = (if\ c = 0\ then\ t - t0\ else\ exp\ (c * t)\ /\ c - exp\ (c * t0)\ /\ c)$
if $t0 \leq t$

for c t::*real*
⟨*proof*⟩

lemma *filtermap-times-pos-at-top*: *filtermap* $((*)\ e)$ *at-top* = *at-top*
if $e > 0$
for e::*real*
⟨*proof*⟩

lemma *exponential-order-additiveI*:
assumes $0 < M$ **and** *eo*: $\forall_F\ t$ *in at-top*. *norm* $(f\ t) \leq K + M * exp\ (c * t)$ **and** $c \geq 0$
obtains M' **where** *exponential-order* M' c f
⟨*proof*⟩

lemma *exponential-order-integral*:
fixes f::*real* ⇒ $'a$::*banach*
assumes I: $\bigwedge t.\ t \geq a \Longrightarrow (f$ *has-integral* $I\ t)\ \{a\ ..\ t\}$
and *eo*: *exponential-order* M c f
and $c > 0$
obtains M' **where** *exponential-order* M' c I
⟨*proof*⟩

lemma *integral-has-vector-derivative-piecewise-continuous*:
fixes f :: *real* ⇒ $'a$::*euclidean-space*— TODO: generalize?
assumes *piecewise-continuous-on* a b D f
shows $\bigwedge x.\ x \in \{a\ ..\ b\} - D \Longrightarrow$
$((\lambda u.$ *integral* $\{a..u\}\ f)$ *has-vector-derivative* $f(x))$ (*at* x *within* $\{a..b\} - D)$
⟨*proof*⟩

lemma *has-derivative-at-split*:
(f *has-derivative* f') (*at* x) ⟷ (f *has-derivative* f') (*at-left* x) ∧ (f *has-derivative* f') (*at-right* x)
for x::$'a$::{*linorder-topology*, *real-normed-vector*}
⟨*proof*⟩

lemma *has-vector-derivative-at-split*:
(f *has-vector-derivative* f') (*at* x) ⟷
(f *has-vector-derivative* f') (*at-left* x) ∧
(f *has-vector-derivative* f') (*at-right* x)
⟨*proof*⟩

lemmas *differentiableI-vector*[*intro*]

lemma *differentiable-at-splitD*:
f *differentiable at-left* x
f *differentiable at-right* x
if f *differentiable* (*at* x)
for x::*real*
⟨*proof*⟩

lemma *integral-differentiable*:
fixes f :: *real* $\Rightarrow$ $'a$::*banach*
assumes *continuous-on* $\{a..b\}$ f
and $x \in \{a..b\}$
shows (λu. *integral* $\{a..u\}$ f) *differentiable at* x *within* $\{a..b\}$
⟨*proof*⟩

theorem *integral-has-vector-derivative-piecewise-continuous'*:
fixes f :: *real* $\Rightarrow$ $'a$::*euclidean-space*— TODO: generalize?
assumes *piecewise-continuous-on* a b D f $a < b$
shows
($\forall x.\ a < x \longrightarrow x < b \longrightarrow x \notin D \longrightarrow$ (λu. *integral* $\{a..u\}$ f) *differentiable at* x) $\wedge$
($\forall x.\ a \leq x \longrightarrow x < b \longrightarrow$ (λt. *integral* $\{a..t\}$ f) *differentiable at-right* x) $\wedge$
($\forall x.\ a < x \longrightarrow x \leq b \longrightarrow$ (λt. *integral* $\{a..t\}$ f) *differentiable at-left* x)
⟨*proof*⟩

lemma *closure* $(-S) \cap$ *closure* S = *frontier* S
⟨*proof*⟩

theorem *integral-time-domain-has-laplace*:
((λt. *integral* $\{0\ ..\ t\}$ f) *has-laplace* L / s) s
if *pc*: $\bigwedge k$. *piecewise-continuous-on* 0 k D f
and *eo*: *exponential-order* M c f
and L: (f *has-laplace* L) s
and s: *Re* $s > c$
and c: $c > 0$
and *TODO*: $D = \{\}$ — TODO: generalize to actual *piecewise-continuous-on*
for f::*real* $\Rightarrow$ *complex*
⟨*proof*⟩

4.4 higher derivatives

definition *nderiv* i f X = ((λf. (λx. *vector-derivative* f (*at* x *within* X)))^^i) f

definition *ndiff* n f X $\longleftrightarrow$ ($\forall i{<}n.\ \forall x \in X$. *nderiv* i f X *differentiable at* x *within* X)

lemma *nderiv-zero*[*simp*]: *nderiv* 0 f X = f
⟨*proof*⟩

lemma *nderiv-Suc*[*simp*]:
nderiv (*Suc* i) f X x = *vector-derivative* (*nderiv* i f X) (*at* x *within* X)
⟨*proof*⟩

lemma *ndiff-zero*[*simp*]: *ndiff* 0 f X
⟨*proof*⟩

lemma *ndiff-Sucs*[*simp*]:
  $ndiff\ (Suc\ i)\ f\ X \longleftrightarrow$
    $(ndiff\ i\ f\ X) \wedge$
    $(\forall x \in X.\ (nderiv\ i\ f\ X)\ differentiable\ (at\ x\ within\ X))$
  ⟨*proof*⟩

theorem *has-laplace-vector-derivative*:
  $((\lambda t.\ vector\text{-}derivative\ f\ (at\ t))\ has\text{-}laplace\ s * L - f0)\ s$
  if L: $(f\ has\text{-}laplace\ L)\ s$
    and f': $\bigwedge t.\ t > 0 \Longrightarrow f\ differentiable\ (at\ t)$
    and $f0$: $(f \longrightarrow f0)\ (at\text{-}right\ 0)$
    and eo: $exponential\text{-}order\ M\ c\ f$
    and cs: $c < Re\ s$
⟨*proof*⟩

lemma *has-laplace-nderiv*:
  $(nderiv\ n\ f\ \{0<..\}\ has\text{-}laplace\ s\hat{}n * L - (\sum i{<}n.\ s\hat{}(n - Suc\ i) * f0\ i))\ s$
  if L: $(f\ has\text{-}laplace\ L)\ s$
    and f': $ndiff\ n\ f\ \{0<..\}$
    and $f0$: $\bigwedge i.\ i < n \Longrightarrow (nderiv\ i\ f\ \{0<..\} \longrightarrow f0\ i)\ (at\text{-}right\ 0)$
    and eo: $\bigwedge i.\ i < n \Longrightarrow exponential\text{-}order\ M\ c\ (nderiv\ i\ f\ \{0<..\})$
    and cs: $c < Re\ s$
  ⟨*proof*⟩

end

5 Lerch Lemma

theory *Lerch-Lemma*
  imports
    HOL−Analysis.Analysis
begin

The main tool to prove uniqueness of the Laplace transform.

lemma *lerch-lemma-real*:
  fixes h::$real \Rightarrow real$
  assumes *h-cont*[*continuous-intros*]: $continuous\text{-}on\ \{0\ ..\ 1\}\ h$
  assumes *int-0*: $\bigwedge n.\ ((\lambda u.\ u\ \hat{}\ n * h\ u)\ has\text{-}integral\ 0)\ \{0\ ..\ 1\}$
  assumes u: $0 \le u\ u \le 1$
  shows $h\ u = 0$
⟨*proof*⟩

lemma *lerch-lemma*:
  fixes h::$real \Rightarrow 'a$::*euclidean-space*
  assumes [*continuous-intros*]: $continuous\text{-}on\ \{0\ ..\ 1\}\ h$
  assumes *int-0*: $\bigwedge n.\ ((\lambda u.\ u\ \hat{}\ n *_R h\ u)\ has\text{-}integral\ 0)\ \{0\ ..\ 1\}$
  assumes u: $0 \le u\ u \le 1$
  shows $h\ u = 0$
⟨*proof*⟩

end

6 Uniqueness of Laplace Transform

theory *Uniqueness*
imports
Existence
Lerch-Lemma
begin

We show uniqueness of the Laplace transform for continuous functions.

lemma *laplace-transform-zero*:— should also work for piecewise continuous
assumes *cont-f*: *continuous-on* $\{0..\}$ f
assumes *eo*: *exponential-order* M a f
assumes *laplace*: $\bigwedge s.$ *Re* $s > a \Longrightarrow$ (f *has-laplace* 0) s
assumes $t \geq 0$
shows $f\ t = 0$
⟨*proof*⟩

lemma *exponential-order-eventually-eq*: *exponential-order* M a f
if *exponential-order* M a g $\bigwedge t.\ t \geq k \Longrightarrow f\ t = g\ t$
⟨*proof*⟩

lemma *exponential-order-mono*:
assumes *eo*: *exponential-order* M a f
assumes $a \leq b$ $M \leq N$
shows *exponential-order* N b f
⟨*proof*⟩

lemma *exponential-order-uminus-iff*:
exponential-order M a $(\lambda x. -f\ x) =$ *exponential-order* M a f
⟨*proof*⟩

lemma *exponential-order-add*:
assumes *exponential-order* M a f *exponential-order* M a g
shows *exponential-order* $(2 * M)$ a $(\lambda x.\ f\ x + g\ x)$
⟨*proof*⟩

theorem *laplace-transform-unique*:
assumes f: $\bigwedge s.$ *Re* $s > a \Longrightarrow$ (f *has-laplace* F) s
assumes g: $\bigwedge s.$ *Re* $s > b \Longrightarrow$ (g *has-laplace* F) s
assumes [*continuous-intros*]: *continuous-on* $\{0..\}$ f
assumes [*continuous-intros*]: *continuous-on* $\{0..\}$ g
assumes *eof*: *exponential-order* M a f
assumes *eog*: *exponential-order* N b g
assumes $t \geq 0$
shows $f\ t = g\ t$
⟨*proof*⟩

end
theory *Laplace-Transform*
imports
Existence
Uniqueness
begin

end

References

[1] A. Rashid and O. Hasan. Formalization of transform methods using HOLăLight. In H. Geuvers, M. England, O. Hasan, F. Rabe, and O. Teschke, editors, *Intelligent Computer Mathematics*, pages 319–332, Cham, 2017. Springer International Publishing.

[2] A. Rashid and O. Hasan. Formalization of Lerch's theorem using HOL Light. *FLAP*, 5(8):1623–1652, 2018.

[3] J. L. Schiff. *The Laplace transform: theory and applications.* Springer New York, 1999.

[4] S. H. Taqdees and O. Hasan. Formalization of Laplace transform using the multivariable calculus theory of HOL-Light. In K. McMillan, A. Middeldorp, and A. Voronkov, editors, *Logic for Programming, Artificial Intelligence, and Reasoning*, pages 744–758, Berlin, Heidelberg, 2013. Springer Berlin Heidelberg.