The Polylogarithm Function

Manuel Eberl

November 9, 2026

Abstract

This entry provides a definition of the *Polylogarithm function*, commonly denoted as $\mathrm{Li}_s(z)$. Here, z is a complex number and s an integer parameter. This function can be defined by the power series expression $\mathrm{Li}_s(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^s}$ for $|z| < 1$ and analytically extended to the entire complex plane, except for a branch cut on $\mathbb{R}_{\geq 1}$.

Several basic properties are also proven, such as the relationship to the Eulerian polynomials via $\mathrm{Li}_{-k}(z) = z(1-z)^{k-1}A_k(z)$ for $k \geq 0$, the derivative formula $\frac{d}{dz}\mathrm{Li}_s(z) = \frac{1}{z}\mathrm{Li}_{s-1}(z)$, the relation to the "normal" logarithm via $\mathrm{Li}_1(z) = -\ln(1-z)$, and the duplication formula $\mathrm{Li}_s(z) + \mathrm{Li}_s(-z) = 2^{1-s}\mathrm{Li}_s(z^2)$.

Contents

1 The Polylogarithm Function **3**
1.1 Definition and basic properties 3
1.2 Special values 5
1.3 Duplication formula 6



Figure 1: Plots of $\mathrm{Li}_s(x)$ for $s = -3, -2, \ldots, 3$ and real inputs $x \in [-2, 1]$

1 The Polylogarithm Function

theory *Polylog*
imports
"HOL-Complex_Analysis.Complex_Analysis"
"Linear_Recurrences.Eulerian_Polynomials"
"HOL-Real_Asymp.Real_Asymp"

begin

1.1 Definition and basic properties

The principal branch of the Polylogarithm function $\mathrm{Li}_s(z)$ is defined as

$$\mathrm{Li}_s(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^s}$$

for $|z| < 1$ and elsewhere by analytic continuation. For integer $s \leq 0$ it is holomorphic except for a pole at $z = 1$. For other values of s it is holomorphic except for a branch cut along the line $[1, \infty)$.

Special values include $\mathrm{Li}_0(z) = \frac{z}{1-z}$ and $\mathrm{Li}_1(z) = -\log(1-z)$.

One could potentially generalise this to arbitrary *s* $\in$ $\mathbb{C}$, but this makes the analytic continuation somewhat more complicated, so we chosed not to do this at this point.

In the following, we define the principal branch of $\mathrm{Li}_s(z)$ for integer s.

definition *polylog :: "int ⇒ complex ⇒ complex"* **where**
"polylog k z =
*(if k ≤ 0 then z * poly (eulerian_poly (nat (-k))) z * (1 - z) powi (k - 1)*
else if z ∈ of_real ‘ {1..} then 0
else (SOME f. f holomorphic_on -of_real‘{1..} ∧
*(∀z∈ball 0 1. f z = (∑n. of_nat (Suc n) powi (-k) * z ^ Suc n))) z)"*

lemma *conv_radius_polylog: "conv_radius (λr. of_nat r powi k :: complex) = 1"*
⟨*proof*⟩

lemma *abs_summable_polylog:*
*"norm z < 1 ⟹ summable (λr. norm (of_nat r powi k * z ^ r :: complex))"*
⟨*proof*⟩

Two very central results that characterise the polylogarithm:

$$\mathrm{Li}_s'(z) = \frac{1}{z}\mathrm{Li}_{s-1}(z) \qquad \text{and} \qquad \mathrm{Li}_s(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^s} \quad \text{for } |z| < 1$$

theorem *has_field_derivative_polylog [derivative_intros]:*
"⋀z. z ∈ (if k ≤ 0 then -{1} else -(of_real ‘ {1..})) ⟹ (polylog k has_field_derivative (if z = 0 then 1 else polylog (k - 1) z / z)) (at z within A)"
and *sums_polylog: "norm z < 1 ⟹ (λn. of_nat (Suc n) powi (-k) * z ^ Suc n) sums polylog k z"*
⟨*proof*⟩

lemma *has_field_derivative_polylog' [derivative_intros]:*
assumes *"(f has_field_derivative f') (at z within A)"*
assumes *"if k ≤ 0 then f z ≠ 1 else Im (f z) ≠ 0 ∨ Re (f z) < 1"*
shows *"((λz. polylog k (f z)) has_field_derivative (if f z = 0 then 1 else polylog (k-1) (f z) / f z) * f') (at z within A)"*
⟨*proof*⟩

lemma *polylog_0 [simp]: "polylog k 0 = 0"*
⟨*proof*⟩

A simple consequence of the derivative formula is the following recurrence for Li_s via a contour integral:

$$\mathrm{Li}_s(z) = \int_0^z \frac{1}{w}\mathrm{Li}_{s-1}(w)\,\mathrm{d}w$$

theorem *polylog_has_contour_integral:*
assumes *"z ∉ complex_of_real ‘ ({..-1} ∪ {1..})"*
shows *"((λw. polylog s w / w) has_contour_integral polylog (s + 1) z) (linepath 0 z)"*
⟨*proof*⟩

lemma *sums_polylog':*
*"norm z < 1 ⟹ k ≠ 0 ⟹ (λn. of_nat n powi - k * z ^ n) sums polylog k z"*
⟨*proof*⟩

lemma *polylog_altdef1:*
*"norm z < 1 ⟹ polylog k z = (∑n. of_nat (Suc n) powi -k * z ^ Suc n)"*
⟨*proof*⟩

lemma *polylog_altdef2:*
*"norm z < 1 ⟹ k ≠ 0 ⟹ polylog k z = (∑n. of_nat n powi -k * z ^ n)"*
⟨*proof*⟩

lemma *polylog_at_pole: "polylog k 1 = 0"*
⟨*proof*⟩

lemma *polylog_at_branch_cut: "x ≥ 1 ⟹ k > 0 ⟹ polylog k (of_real x) = 0"*
⟨*proof*⟩

lemma *holomorphic_on_polylog [holomorphic_intros]:*
assumes *"A ⊆ (if k ≤ 0 then -{1} else -of_real ` {1..})"*
shows *"polylog k holomorphic_on A"*
⟨*proof*⟩

lemmas *holomorphic_on_polylog' [holomorphic_intros] = holomorphic_on_compose_gen [OF _ holomorphic_on_polylog[OF order.refl], unfolded o_def]*

lemma *analytic_on_polylog [analytic_intros]:*
assumes *"A ⊆ (if k ≤ 0 then -{1} else -of_real ` {1..})"*
shows *"polylog k analytic_on A"*
⟨*proof*⟩

lemmas *analytic_on_polylog' [analytic_intros] = analytic_on_compose_gen [OF _ analytic_on_polylog[OF order.refl], unfolded o_def]*

lemma *continuous_on_polylog [analytic_intros]:*
assumes *"A ⊆ (if k ≤ 0 then -{1} else -of_real ` {1..})"*
shows *"continuous_on A (polylog k)"*
⟨*proof*⟩

lemmas *continuous_on_polylog' [continuous_intros] = continuous_on_compose2 [OF continuous_on_polylog [OF order.refl]]*

1.2 Special values

lemma *polylog_neg_int_left:*
*"k < 0 ⟹ polylog k z = z * poly (eulerian_poly (nat (-k))) z * (1 - z) powi (k - 1)"*
⟨*proof*⟩

lemma *polylog_0_left: "polylog 0 z = z / (1 - z)"*
⟨*proof*⟩

lemma *polylog_neg1_left: "polylog (-1) x = x / (1 - x) ^ 2"*
⟨*proof*⟩

lemma *polylog_neg2_left: "polylog (-2) x = x * (1 + x) / (1 - x) ^ 3"*
⟨*proof*⟩

lemma *polylog_neg3_left: "polylog (-3) x = x * (1 + 4 * x + x²) / (1 - x) ^ 4"*
⟨*proof*⟩

lemma *polylog_1:*
  assumes *"z ∉ of_real ‘ {1..}"*
  shows *"polylog 1 z = -ln (1 - z)"*
⟨*proof*⟩

lemma *is_pole_polylog_1:*
  assumes *"k ≤ 0"*
  shows *"is_pole (polylog k) 1"*
⟨*proof*⟩

lemma *zorder_polylog_1:*
  assumes *"k ≤ 0"*
  shows *"zorder (polylog k) 1 = k - 1"*
⟨*proof*⟩

lemma *isolated_singularity_polylog_1:*
  assumes *"k ≤ 0"*
  shows *"isolated_singularity_at (polylog k) 1"*
  ⟨*proof*⟩

lemma *not_essential_polylog_1:*
  assumes *"k ≤ 0"*
  shows *"not_essential (polylog k) 1"*
  ⟨*proof*⟩

lemma *polylog_meromorphic_on [meromorphic_intros]:*
  assumes *"k ≤ 0"*
  shows *"polylog k meromorphic_on {1}"*
  ⟨*proof*⟩

1.3 Duplication formula

Lastly, we prove the following duplication formula that the polylogarithm satisfies:

$$\mathrm{Li}_s(z) + \mathrm{Li}_s(-z) = 2^{1-s}\mathrm{Li}_s(z^2)$$

The proof is a relatively simple manipulation of infinite sum that defines $\mathrm{Li}_s(z)$ for $|z| < 1$, followed by analytic continuation to its full domain.

theorem *polylog_duplication:*
  assumes *"if s ≤ 0 then z ∉ {-1, 1} else z ∉ complex_of_real ‘ ({..-1} ∪ {1..})"*
  shows *"polylog s z + polylog s (-z) = 2 powi (1 - s) * polylog s (z²)"*
⟨*proof*⟩

end

References

[1] J. Mason and D. Handscomb. *Chebyshev Polynomials*. CRC Press, 2002.