Ramsey's Theorem

Tom Ridge

June 17, 2024

Abstract

The infinite form of Ramsey's Theorem is proved following Boolos and Jeffrey, Chapter 26.

Contents

1 Ramsey's Theorem

theory *Ramsey*
imports *Main HOL−Library.Infinite-Set HOL−Library.Ramsey*

begin

Please note: this entire development has been updated and incorporated into *HOL−Library.Ramsey* above. Below, some of the results of the original development are linked to their current versions elsewhere in the Isabelle libraries.

1.1 Library lemmas

lemma *infinite-inj-infinite-image*: *infinite* $Z \Longrightarrow$ *inj-on* $f\ Z \Longrightarrow$ *infinite* $(f \text{ ` } Z)$
⟨*proof*⟩

lemma *infinite-dom-finite-rng*: [| *infinite* A; *finite* $(f \text{ ` } A)$ |] ==> $\exists\, b \in f \text{ ` } A.$ *infinite* $\{a : A.\ f\ a = b\}$
⟨*proof*⟩

lemma *infinite-mem*: *infinite* $X \Longrightarrow \exists\, x.\ x \in X$
⟨*proof*⟩

lemma *not-empty-least*: $(Y{::}nat\ set) \neq \{\} \Longrightarrow \exists\, m.\ m \in Y \wedge (\forall\, m'.\ m' \in Y \longrightarrow m \leq m')$
⟨*proof*⟩

1.2 Dependent Choice Variant

lemma *dc*:
assumes *trans*: $trans\ r$
and *P0*: $P\ x0$
and *Pstep*: $\bigwedge x.\ P\ x \Longrightarrow \exists\, y.\ P\ y \wedge (x,\ y) \in r$
obtains $f :: nat \Rightarrow {'a}$ **where** $\bigwedge n.\ P\ (f\ n)$ **and** $\bigwedge n\ m.\ n < m \Longrightarrow (f\ n,\ f\ m) \in r$
⟨*proof*⟩

1.3 Ramsey's theorem

lemma *ramsey*: $\forall\, (s{::}nat)\ (r{::}nat)\ (YY{::}{'a}\ set)\ (f{::}{'a}\ set \Rightarrow nat).$
$infinite\ YY$
$\wedge\ (\forall\, X.\ X \subseteq YY \wedge finite\ X \wedge card\ X = r \longrightarrow f\ X < s)$
$\longrightarrow (\exists\, Y'\ t'.$
$Y' \subseteq YY$
$\wedge\ infinite\ Y'$
$\wedge\ t' < s$
$\wedge\ (\forall\, X.\ X \subseteq Y' \wedge finite\ X \wedge card\ X = r \longrightarrow f\ X = t'))$
⟨*proof*⟩

end